

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Microplastics and their associated organic pollutants from the coastal waters of the central Adriatic Sea (Italy): Investigation of adipogenic effects in vitro**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1755254> since 2020-12-18T18:40:22Z

*Published version:*

DOI:10.1016/j.chemosphere.2020.128090

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

# Journal Pre-proof

Microplastics and their associated organic pollutants from the coastal waters of the central Adriatic Sea (Italy): investigation of adipogenic effects *in vitro*

Martina Capriotti, Paolo Cocci, Luca Bracchetti, Erika Cottone, Rosaria Scandiffio, Giovanni Caprioli, Gianni Sagratini, Gilberto Mosconi, Patrizia Bovolín, Francesco Alessandro Palermo

PII: S0045-6535(20)32285-2

DOI: <https://doi.org/10.1016/j.chemosphere.2020.128090>

Reference: CHEM 128090

To appear in: *ECSN*

Received Date: 25 May 2020

Revised Date: 28 July 2020

Accepted Date: 19 August 2020



Please cite this article as: Capriotti, M., Cocci, P., Bracchetti, L., Cottone, E., Scandiffio, R., Caprioli, G., Sagratini, G., Mosconi, G., Bovolín, P., Palermo, F.A., Microplastics and their associated organic pollutants from the coastal waters of the central Adriatic Sea (Italy): investigation of adipogenic effects *in vitro*, *Chemosphere*, <https://doi.org/10.1016/j.chemosphere.2020.128090>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Elsevier Ltd. All rights reserved.

**CRedit author statement**

Martina Capriotti: Investigation, Resources, Writing- Original draft; Paolo Cocci: Methodology, Investigation, Writing- Original draft; Luca Bracchetti: Investigation; Erika Cottone: Methodology, Writing- Original draft; Rosaria Scandiffio: Investigation, Data curation; Giovanni Caprioli: Investigation, Methodology; Gianni Sagratini: Investigation, Methodology; Gilberto Mosconi: Data curation; Patrizia Bovolín: Formal analysis, Writing- Original draft; Francesco Alessandro Palermo: Conceptualization, Data curation, Supervision, Writing - Original Draft, Writing- Reviewing and Editing.

**Microplastics and their associated organic pollutants from the coastal waters of the central Adriatic Sea (Italy): investigation of adipogenic effects *in vitro***

Martina Capriotti<sup>1,2</sup>, Paolo Cocci<sup>2</sup>, Luca Bracchetti<sup>2</sup>, Erika Cottone<sup>3</sup>, Rosaria Scandiffio<sup>3</sup>, Giovanni Caprioli<sup>4</sup>, Gianni Sagratini<sup>4</sup>, Gilberto Mosconi<sup>2</sup>, Patrizia Bovolín<sup>4</sup>, Francesco Alessandro Palermo<sup>2\*</sup>

1. Department of Marine Sciences, University of Connecticut, 1080 Shennecossett Rd, Groton (CT) USA

2. School of Biosciences and Veterinary Medicine, University of Camerino, Via Gentile III da Varano, 62032 Camerino (MC) ITALY

3. Department of Life Sciences and Systems Biology, University of Turin, Via Accademia Albertina 13, 10123 Torino (TO) ITALY

4. School of Pharmacy, University of Camerino, Via Madonna delle Carceri 9, 62032 Camerino (MC) ITALY

**\*Correspondence may be addressed to:**

Francesco A. Palermo

School of Biosciences and Veterinary Medicine, University of Camerino,

Via Gentile III Da Varano, I-62032 Camerino (MC), Italy

Tel. +39 0737 404920

Fax +39 0737 404901

e-mail francesco.palermo@unicam.it

**Abstract**

Even though microplastic (MP) pollution in aquatic environment is nowadays widely studied, a huge gap of knowledge exists on their actual biological effects. In this study we first reported environmental baseline data on the occurrence and characterization of floating MPs in Italian coastal waters of the Central Adriatic Sea by using a standardized monitoring protocol. Further, we analyzed the concentrations of MP-associated chemicals and evaluated their potential adipogenic effects using 3T3-L1 preadipocytes. MPs were found in each sampling stations showing the highest abundance ( $1.88 \pm 1.78$  items/m<sup>3</sup>) in the sites more distant from the coast with fragments as the most common shape category. All targeted organic pollutants (*i.e.* polychlorinated biphenyls - PCBs, polycyclic aromatic hydrocarbons -PAHs, organophosphorus - OP, and organochlorine - OC pesticides) have been detected on the surface of the collected MPs. The highest concentrations of PAHs were found on MPs from inshore (*i.e.* <1.5 NM) surface waters with low-ring PAHs as dominant components. Similarly, MPs from inshore waters had higher  $\Sigma$ PCB concentrations (64.72 ng/g plastic) than those found in offshore (*i.e.* >6 NM) waters (10.37 ng/g plastic). Among pesticides, all measured OPs were detected in each sample analyzed with pirimiphos-methyl as the most representative compound. For OCs, the sum of all concentrations of congeners was higher in coastal with respect to offshore waters. Moreover, *in vitro* 3T3-L1 screening of MP extracts indicated potential metabolic effects resulting in both adipogenesis and lipid uptake/ storage.

**Keywords:** Adriatic Sea; Microplastic; Biomonitoring; Endocrine disruptors; Metabolic disruptors; Adipocytes

## 1. Introduction

Microplastics (MPs,  $\geq 0.001\text{mm}$  and  $< 5\text{mm}$ ) (Shim et al., 2018) are a well known global issue for marine and coastal ecosystems (Browne et al., 2011; UNEP, 2016) due to their ubiquitous presence throughout the marine environment compartments, as surface and water column, sediments, and biota. In this regard, the Mediterranean Sea with a mean density of floating microplastics of about 100,000 items/ $\text{km}^2$  represents one of the basin most affected by MP litter (Fossi et al., 2012). Within the Mediterranean Sea, however, very few studies have investigated the occurrence of floating plastic debris in the Adriatic Sea that, due to its features (*e.g.* semi-closed conformation, oceanographic conditions and heavy anthropogenic pressure), can be considered a plastic pollution hot spot (Gomiero et al., 2018). In this regard, recent data on the Mediterranean Sea (partially including the Adriatic area) reported a MP range of 0.4 - 1.8 items/ $\text{m}^3$  with polyethylene (PE) as the most common polymer composition types (Suaria et al., 2016). Similarly, a clear prevalence of MPs (65.1% of sampled debris) was demonstrated in marine sediments from the Central Adriatic (Munari et al., 2017). MPs were also found in a wide range of marine species collected along the Adriatic Sea showing highest concentrations (*i.e.* 1.0–1.7 items/specimens) in fish (Avio et al., 2015; Pellini et al., 2018). These findings indicate that small size debris are available to marine vertebrates and invertebrates, resulting in both physical and chemical impacts (Fossi et al., 2012; Fossi et al., 2014; Gall and Thompson, 2015). MPs can be translocated to the lymphatic and/or circulatory system, and accumulated in organs as the digestive and the respiratory tracts (Barnes et al., 2009; Cole and Galloway, 2015; Mazurais et al., 2015; Besley et al., 2017; Lusher et al., 2017a; Lusher et al., 2017b).

Nowadays the research on the ecological impact of microplastics is not limited to evaluate their accumulation in the marine environment and consequently in living organisms, but is now moving on to investigate the role of MPs as potential vehicles of a wide range of toxic chemicals (Teuten et al., 2007; Engler, 2012). Recent data indicate that hydrophobic organic contaminants (HOCs), including endocrine disrupting chemicals (EDCs), have been detected on the surface of MPs (Mato

et al., 2001; Ogata et al., 2009; Rios et al., 2010; Hirai et al., 2011; Rochman et al., 2014a). Indeed, due to the physical and chemical properties of plastic polymers, MPs can quickly accumulate and concentrate HOCs present in the surrounding water. It should also be taken into account that MP-associated HOCs have to be added to the chemicals already included as additives during plastic manufacturing (*e.g.* bisphenol A or nonylphenol) (Ogata et al., 2009; Hirai et al., 2011; Engler, 2012; Mai et al., 2018; Alimba and Faggio, 2019; Chen et al., 2019), making the potential of chemicals release from MPs to the marine environment and wildlife particularly harmful. In addition, ingestion of MPs by aquatic animals may result in increased bioavailability of HOCs, bioaccumulation and both potential toxicity and transfer along the trophic web (Avio et al., 2015; Batel et al., 2016).

It has been demonstrated that among HOCs, EDCs are commonly found on MPs, including abrasion beads, and easily released from plastic debris due to lower partition coefficients between plastic and water (Liu et al., 2016). EDCs accumulation in tissues of marine organisms, following their MP-mediated release, can thus exert adverse effects ranging from disruption of endocrine-based reproductive pathways to metabolic alterations (Franzellitti et al., 2019). In this last regard, a group of endocrine disruptors, named obesogens or metabolic disruptors, have been found to affect adipogenesis by perturbing peroxisome proliferator activated receptor (PPAR) signaling pathways (Grün and Blumberg, 2009). Recently, the obesogenic potential of these pollutants were assayed *in vitro* using 3T3-L1 adipocytes and primary cultures of sea bream hepatocytes (Cocci et al., 2017; Pomatto et al., 2018).

Thus, our first objective in this study was to collect environmental baseline data on the occurrence and characterization of floating MPs in Italian coastal waters of the Central Adriatic Sea by using a standardized monitoring protocol. Second, we aimed to examine the concentrations of chemicals adsorbed on MPs sampled from the same study area and to test their adverse effect by *in vitro* bioassays. In particular, we used 3T3-L1 preadipocytes to investigate the possible adipogenic

effects of these plastic extracts in order to provide preliminary insights on potential ecotoxicological risk of MPs as vehicles of chemical pollutants.

## 2. Materials and Methods

### 2.1 Sites and water surface sampling

Samples were collected inshore and offshore along the western (Italian) coasts of the Central Adriatic Sea (Province of Ascoli Piceno, Marche Region - Italy). The sampling operations were conducted during summer season 2018, following the microplastics monitoring methods applied by the Italian Ministry of the Environment and Land and Sea protection (Minambiente, 2018) and the requirements indicated by the ministerial protocol within the MSFD monitoring program (MSFD, 2013). Briefly, transects were outlined perpendicular to the coast line along the municipalities of San Benedetto del Tronto (SBT) and Grottammare (GRT), respectively (Figure 1). Three sampling stations were located at different distance from the coast (0.5, 1.5 and 6 nautical Miles, NM) along each transect. An additional sampling station was located inside the San Benedetto del Tronto harbor. Floating samples were collected from surface sea water, using a manta trawl with mesh size of 300  $\mu\text{m}$  (30 x 15 x 200 cm) equipped with a mechanical flow meter (Hydro-Bios). The manta net was towed on the water surface for 20 minutes at 2 knots and was kept at a distance far from the boat to avoid turbulence by the waking of the ship. Samples were rinsed from the outside to the end of the net, placed in glass containers and immediately stored at 4 °C until the sorting using a stereomicroscope. To prevent external contamination during the analysis, the laboratory procedures were performed according to Baini et al. (2018).






Figure 1. Map of the study area. Sampling stations were located at different distance from the coast (0.5, 1.5 and 6 nautical Miles, NM) along each transect.

## 2.2 Microplastic identification and count

Samples were dried at room temperature and weighed in OHAUS Explorer analytical balance. For microplastics identification, samples were observed under a stereomicroscope (Carl Zeiss Stemi<sup>TM</sup>) and images have been examined by a USB Camera (Optika B Series) using the Optika ProView software. Plastic particles were sorted based on their colour (blue, red, black, white, transparent, green, other colour), size (<5, 5–25, >25 mm) and shape (spherical, filament, fragment, sheet, other shape) according to the MSFD guidelines and Imhof et al. (2017). As reported by Baini et al. (2018), the number of microplastic was normalized to the total water volume filtered (V) and expressed as items/m<sup>3</sup>.

### 2.3 Determination of PAHs, PCBs, Organophosphorus and organochlorine pesticides

Due to limited amount of particles to be provided for both bioassay and chemical analysis, MPs collected within the study area were pooled according to distance from the coast in order to obtain 3 type of samples: offshore waters (*i.e.* 6 NM), inshore waters (*i.e.* <1.5 NM, corresponding to local areas of touristic and commercial traffics) and the harbor of San Benedetto del Tronto as expected polluted area.

Hydrophobic organic contaminants were isolated from marine debris following the protocol described by Chen et al. (2019) with slight modifications. Briefly, 4 mL of dichloromethane was added to each microplastic sample (400 mg) and incubated for 30 min at room temperature, vortexing every 10 minutes. The procedure was repeated for 3 times and finally the supernatant was evaporated to 500 µl using a Rotavapor R-300 (Buchi). The PCB mix standard (PCB 28, PCB 52, PCB 95, PCB 99, PCB 101, PCB 105, PCB 110, PCB 118, PCB 138, PCB 146, PCB 149, PCB 151, PCB 153, PCB 170, PCB 177, PCB 180, PCB 183, PCB 187) at a concentration of 10 mg/L in iso-octane, the organochlorine pesticides OC1 (six congeners, *i.e.*  $\alpha$ -HCH,  $\beta$ -HCH,  $\gamma$ -HCH, HCB, heptachlor and heptachlor epoxide) mix standard at a concentration of 10 mg/L in cyclohexane and the organochlorine pesticides OC2 (six congeners, *i.e.* 2,4-DDT, 4,4-DDT, 2,4-DDE, 4,4-DDE, 2,4-DDD, 4,4-DDD) mix standard at a concentration of 10 mg/L in cyclohexane were supplied by Dr. Ehrenstorfer (Ausburg, Germany). Standard working solutions at various concentrations were prepared daily by appropriate dilution of the stock solutions with hexane. The organophosphorus pesticides (four congeners, *i.e.* chlorfenvinphos, chlorpyrifos, malathion, pirimiphos-methyl) mix standard were supplied by Fluka (Milano, Italy). Individual stock solutions of organophosphorus pesticides at concentrations of 1000 mg/L were prepared by dissolving pure standard compounds in HPLC grade methanol and then stored in glass-stoppered bottles at 4 °C. Afterwards standard working solutions at various concentrations were prepared daily by appropriate dilution of the stock solution with methanol. A gas chromatograph/mass selective detector (GC/MSD) (Hewlett Packard, Palo Alto, CA, USA; HP-6890 with HP 5973) was used. Separation was performed on an HP 5 MSI

column (30 m X 0.25 mm X 0.25  $\mu$ m film thickness). An HP Chem workstation was used with the GC/MS system. All injections were splitless and the volume was 1  $\mu$ l. The flow rate (He) was 0.8 mL/min. The injector temperature was 270 °C. The column temperature program used for PCB and for the second group of organochlorine pesticides, *i.e.* dichlorodiphenyltrichloroethane (DDT) and its metabolites (OC 2 subgroup) analyses was from 60 °C (3 min) to 190 °C at 8 °C min<sup>-1</sup>, from 190 °C (18 min) to 300 °C at 15 °C min<sup>-1</sup>, then at 300 °C for 1 min. The column temperature programme used for the first group of organochlorine pesticides, *i.e.*  $\alpha$ -HCH,  $\beta$ -HCH,  $\gamma$ -HCH, HCB, heptachlor and heptachlor epoxide (all included in the OC 1 subgroup) was from 60 °C (3 min) to 190 °C at 8 °C min<sup>-1</sup>, from 190 °C (13 min) to 300 °C at 15 °C min<sup>-1</sup>, then at 300 °C for 1 min. The column temperature programme used for organophosphorus pesticides analyses (chlorfenvinphos, chlorpyrifos, malathion, pirimiphos-methyl) was from 50 °C (5 min) to 320 °C at 15 °C min<sup>-1</sup>, then at 320 °C for 4 min. Data were acquired in the electron impact (EI) mode (70 eV) using the selected ion monitoring (SIM) mode. For this purpose, a gas chromatograph/mass selective detector (GC/MSD) (Hewlett Packard. Palo Alto. CA. USA; HP-6890 with HP 5973) was used. Extracts were analyzed for concentrations of 15 of the most environmentally relevant PAHs (Naphthalene, Acenaphthene, Fluorene, Chrysene, Phenanthrene, Fluoranthene, Anthracene, Pyrene, Benzo[a]anthracene, Benzo[b]fluoranthene, Benzo[k]fluoranthene, Benzo[a]pyrene, Dibenzo[a,h]anthracene, Benzo[g,h,i]perylene and Indeno[1,2,3-c,d]pyrene). The detailed methodology for separation and quantification of target analytes was published previously (see “Supplementary data” of Cocci et al., 2017).

#### 2.4 3T3-L1 cell culture and quantification of adipocyte lipid accumulation

3T3-L1 preadipocytes (ATCC<sup>®</sup> CL-173<sup>™</sup>; Lot No 70009858, ATCC, Manassas, VA, USA) were cultured in high-glucose (4.5 g/L) Dulbecco’s modified Eagle’s medium (DMEM) supplemented with 10% bovine calf serum, 2 mM L-glutamine, 50 IU/mL penicillin, and 50  $\mu$ g/mL streptomycin (Sigma Aldrich, St. Louis, MO, USA).

191 For experiments,  $5 \times 10^3$  cells/well were seeded in 96-black well clear bottom plates (Greiner Bio-  
192 One, Frickenhausen, Germany). Two days after reaching confluence (day 0), cells were incubated  
193 with differentiation medium (MDI; DMEM containing 10% fetal bovine serum, 1  $\mu\text{g/mL}$  insulin,  
194 0.5 mM isobutylmethylxanthine), containing different concentrations of the various microplastic  
195 organic extracts. In particular, inshore, offshore and harbor microplastic extracts were dried and  
196 redissolved in 30  $\mu\text{l}$  DMSO; then scalar dilutions, ranging from  $10^{-3}$  to  $10^{-6}$ , were added to the MDI  
197 used for adipocyte differentiation. Two days later (day 2), MDI medium was replaced with  
198 maintenance medium (MM; DMEM 10% FBS, 1  $\mu\text{g/mL}$  insulin), always containing the various  
199 dilutions of microplastic extracts. Negative controls (solvent only) were set up using MDI and MM  
200 added with 0.1% DMSO, corresponding to the maximum solvent concentration present in  
201 microplastic extract dilutions; as a positive control, 100 nM Rosiglitazone (Sigma Aldrich) was  
202 added to MDI and MM. Fresh medium was provided every two days; experiments ended after 9  
203 days from the beginning of the differentiation (day 9). Lipid accumulation was quantified by using  
204 AdipoRed<sup>TM</sup> assay reagent (Lonza, Walkersville, MD, USA), while the DNA content (that  
205 correlates with cell number) was estimated by NucBlue<sup>TM</sup> staining (Invitrogen, Carlsbad, CA,  
206 USA). Briefly, medium was removed from 3T3-L1 cultures and cells were rinsed with PBS,  
207 subsequently replaced with a dye mixture containing AdipoRed<sup>TM</sup> and NucBlue<sup>TM</sup> assay reagents  
208 diluted in PBS (25  $\mu\text{l}$  and 1 drop, respectively, per mL of PBS). After 40 min of incubation at room  
209 temperature in the dark, fluorescence was measured with Filtermax F5 microplate reader  
210 (Molecular Devices, Sunnyvale, CA, USA) with excitation at 485 nm and emission at 535 nm for  
211 AdipoRed<sup>TM</sup> and excitation at 360 nm and emission at 460 nm for NucBlue<sup>TM</sup> quantification.  
212 Experiments were repeated three times (four wells for each condition), using cells at different  
213 passage numbers (p3-p5). Data were referred as % of the control (0.1% DMSO) condition (set =  
214 100%).

## 2.5 Statistical analysis

Data are expressed as mean  $\pm$  standard error of the mean (SEM). Statistical analysis was performed using ANOVA (one-way analysis of variance) followed by Bonferroni's multiple comparison test. Differences with  $p < 0.05$  were considered statistically significant.

## 3. Results

### 3.1 Microplastics in water surface samples

Particles isolated from surface seawater were visually counted and sorted based on their dimension, shape and color (Table 1). Among the total particles sampled, microplastics ( $< 5\text{mm}$ ) constituted the most representative size class ( $n=401.08$ , 95%), and were detected in each transect representing the totality of plastic litter (100%) in GRT1, GRT2 and SBT2 stations and showing the highest abundance in the sites more distant from the coast (SBT3:  $1.88 \pm 1.78$  and GTR3:  $3.42 \pm 2.28$  items/ $\text{m}^3$ ) or inside the San Benedetto del Tronto harbor ( $0.91 \pm 0.58$  items/ $\text{m}^3$ ). Marine debris larger than 5 mm were recorded in harbor (16%), open waters (SBT3, 6% and GRT3, 12%), and also at 0.5 NM along the SBT transect (2%).

*Table 1. Particles abundance (items/ $\text{m}^3$ ) and size (mm) in water surface samples collected in three different sampling sites; mean values  $\pm$  s.d. for each transect have been reported.*

Sampling site	Distance (nM)	Items/ $\text{m}^3$	Micro- ( $< 5$ mm)	Meso- (5 to 25 mm)	Macro- ( $> 25$ mm)
GRT1	0.5	$0.17 \pm 0.02$	$12.33 \pm 2.51$ (100%)	-	-
GRT2	1.5	$0.11 \pm 0.05$	$9.34 \pm 4.01$ (100%)	-	-
GRT3	6	$3.42 \pm 2.28$	$240.33 \pm 160.55$ (88%)	$38.33 \pm 33.29$ (10%)	$1.67 \pm 1.53$ (2%)

<i>SBT1</i>	0.5	$0.17 \pm 0.11$	$15.75 \pm 10.59$ (98%)	$0.50 \pm 1.00$ (2%)	-	235
<i>SBT2</i>	1.5	$0.04 \pm 0.01$	$3.00 \pm 0.00$ (100%)	-	-	236
<i>SBT3</i>	6	$1.88 \pm 1.78$	$66.00 \pm 8.54$ (94%)	$3.33 \pm 0.58$ (5%)	$1.0 \pm 1.00$ (1%)	237
<i>Harbor</i>		$0.91 \pm 0.57$	$54.33 \pm 24.11$ (84%)	$11.67 \pm 15.30$ (11%)	$4.33 \pm 4.51$ (5%)	238
						239

240

241 Analyzing the overall study area by combining the values measured at two sampling stations at the  
242 same distance (along both transects), 75% of the total MPs were collected from sampling sites at 6  
243 NM, 3% at 1.5 NM, 8% at 0.5 NM from the coastline and finally the 13% inside the harbor.  
244 Fragments were the most common shape category observed in each sampling station (Figure 2a),  
245 reaching the 74.7% (n= 28.4) of all the MP types sampled in coastal sites (<1.5 NM). Films were  
246 the second most abundant group of the plastic items collected with the highest number (n= 23.3,  
247 42.9%) recorded in SBT harbor. On the other hand, spherical particles were commonly found in  
248 offshore waters (6 NM) showing a total number of 54 (25%, Figure 2a).

249 Analysis of MP color pattern revealed that transparent and white items accounted for 37% and 44%  
250 of the particles collected respectively (Figure 2b).

251




Figure 2. Sampling station-related abundance % of microplastics isolated from water surface samples in relation to shape (a) and color (b).

### 3.2 Concentrations of pollutants on inshore and offshore microplastic samples

MP extracts were analyzed for PCBs, PAHs, OPs, and OCs (Table 2).

*Table 2. Concentration (ng/g plastic) of Polychlorinated biphenyls (PCBs), organophosphorus (OPs), organochlorine pesticides (OCs), and polycyclic aromatic hydrocarbon (PAH) congeners measured on microplastics from inshore (i.e. <1.5 NM), offshore waters (i.e. 6 NM), and harbor.*

POPs	Inshore (ng/g plastic)	Offshore (ng/g plastic)	Harbor (ng/g plastic)
PCB 28	n.d.	n.d.	n.d.
PCB 52	27.91	0.70	12.22
PCB 95	3.58	0.17	5.85
PCB 99	0.60	n.d.	n.d.
PCB 101	n.d.	0.08	0.52
PCB 105	n.d.	n.d.	n.d.
PCB 110	n.d.	0.08	0.34
PCB 118	n.d.	n.d.	n.d.
PCB 138	28.51	0.54	16.01
PCB 146	n.d.	n.d.	n.d.
PCB 149	5.07	0.29	2.84
PCB 151	n.d.	n.d.	n.d.
PCB 153	n.d.	n.d.	n.d.
PCB 170	n.d.	n.d.	n.d.
PCB 177	n.d.	n.d.	n.d.
PCB 180	n.d.	n.d.	n.d.
PCB 183	n.d.	n.d.	n.d.
PCB 187	n.d.	n.d.	n.d.
$\Sigma$ PCBs	65.67	1.86	37.78
Chlorfenvinphos	6.57	0.14	1.98
Chlorpyrifos	45.37	0.77	32.19
Malathion	17.31	0.22	7.23
Pirimiphos-methyl	78.36	3.59	32.70
$\Sigma$ OPs	147.61	4.72	74.10
$\alpha$ -HCH	95.22	0.84	27.02
$\beta$ -HCH	0.37	0.02	0.86
$\gamma$ -HCH	3.58	0.07	1.98
HCB	2.09	0.10	10.84
Heptachlor	n.d.	n.d.	n.d.
Heptachlor epoxide	n.d.	n.d.	n.d.
$\Sigma$ OC1s	101.26	1.03	40.70
2,4-DDE	10.15	0.16	2.41
4,4-DDE	8.66	0.19	n.d.



2,4-DDD	n.d.	1.32	8.00
4,4-DDD	15.52	n.d.	67.38
2,4-DDT	n.d.	0.59	n.d.
4,4-DDT	n.d.	2.79	n.d.
$\Sigma$ OC2s	34.33	5.05	77.79
Naphthalene	9.55	n.d.	6.71
Acenaphthene	n.d.	n.d.	n.d.
2-Bromonaphthalene	46.27	2.46	40.36
Acenaphthylene	109.40	1.27	100.00
Fluorene	n.d.	n.d.	n.d.
Phenanthrene	n.d.	n.d.	2.58
Anthracene	64.18	1.54	25.04
Fluoranthene	16.87	0.57	n.d.
Pyrene	n.d.	1.48	35.13
Benzo(a)Anthracene	67.31	2.64	40.36
Chrysene	81.64	2.28	47.93
Benzo(b)Fluoranthene	n.d.	n.d.	n.d.
Benzo(a)Pyrene	n.d.	n.d.	7.66
Indeno(1,2,3-cd)Pyrene	n.d.	n.d.	n.d.
DiBenz(a,h)Anthracene	n.d.	4.40	12.91
Benzo(g,h,i)Perylene	12.84	2.34	41.48
$\Sigma$ low-ring PAHs	229.40	5.27	174.69
$\Sigma$ high-ring PAHs	178.66	13.71	185.47
$\Sigma$ Total PAHs	408.06	18.98	360.16

---


n.d.: not detectable.

The highest concentrations of PAHs were found in microplastics collected from both inshore and harbor surface waters (Figure 3) with low-ring PAHs as dominant components (Table 2). On the contrary, high-ring PAHs were most abundant in offshore samples (Table 2).

Among the sixteen PCB congeners analysed, only seven, namely PCB 52, PCB 95, PCB 99, PCB 101, PCB 110, PCB 138 and PCB 149, were detected in the different samples (Table 2). The congeners 52, 95, 138 and 149 were found in MPs from each sampling area. If congeners were considered altogether, inshore waters had the highest  $\Sigma$ PCB concentrations (64.72 ng/g plastic) followed by SBT harbor (37.86 ng/g plastic). On the contrary, the lowest  $\Sigma$ PCBs was found in MPs from offshore waters (10.37 ng/g plastic) (Figure 3).

All measured OPs were detected in each sample analyzed with pirimiphos-methyl as the most representative compound. Generally, the highest  $\Sigma$ OP concentrations were found in samples from

276 inshore waters (Figure 3). For OC1 and OC2, the sum of all concentrations of congeners were  
 277 higher in coastal waters with respect to offshore waters (Figure 3).



278

279 *Figure 3. Microplastic-associated PAH, PCB, OP, OC1 and OC2 patterns across sampling sites.*


280

281

### 282 3.3 Effects of microplastic extracts on 3T3-L1 adipocyte differentiation

283 The effect of MP extracts was assayed on the murine 3T3-L1 preadipocyte cell line, a commonly  
 284 used cell model for adipose cell biology research. Since adipogenic effects can be exerted by  
 285 increasing intracellular lipids (adipocytes hypertrophy) and/or adipocytes number (adipocytes  
 286 hyperplasia), we assayed both triglyceride accumulation (AdipoRed<sup>TM</sup> assay) and total cell number  
 287 (NucBlue<sup>TM</sup> staining, measuring DNA content). Confluent preadipocytes, cultured in 96-well  
 288 plates, were induced to start adipogenic differentiation and were treated throughout the  
 289 differentiation period (9 days) with solvent only (0.1% DMSO; negative control) or with different  
 290 concentrations of microplastic extracts (serial dilutions ranging from  $10^{-3}$  to  $10^{-6}$ ); Rosiglitazone, a  
 291 well-known agonist of the adipogenesis master regulator PPAR $\gamma$ , was used as a positive control. All  
 292 the tested concentrations of microplastic extracts did not exert cytotoxic effects, as demonstrated by

cell evaluation under the microscope and by NucBlue staining quantification, showing no difference in DNA content, and thus in cell number, in microplastic extract-exposed cells versus control adipocytes (Figure 4c). Interestingly, harbor, inshore and offshore microplastic extracts, at all the concentrations used, did induce an increase in triglyceride accumulation, in respect to control cells (Figure 4a, b). Differently from Rosiglitazone treatment, exposure to microplastic extracts induced no changes in total cell number (Figure 4c), thus indicating that microplastic extracts did not induce adipocytes hyperplasia. On the other hand, triglyceride accumulation per cell (i.e. triglyceride accumulation normalized for DNA content) resulted statistically increased after microplastic extracts exposure (Figure 4d), thus suggesting an induction of adipocytes hypertrophy.



**Figure 4.** (a) Representative micrographs of AdipoRed (red) and NucBlue (blue) staining of undifferentiated 3T3-L1 preadipocytes (Undiff.) and adipocytes after 9 days of differentiation in the presence of vehicle only (0.1% DMSO; Control), Rosiglitazone, or extracts obtained from microplastics collected at the different sampling stations (Harbor, Inshore, Offshore). Scale bar: 50 μm. (b) Graph summarizing AdipoRed staining experiments to assess lipid accumulation in differentiated 3T3-L1 adipocytes treated with 0.1% DMSO solvent (control) or cells treated with 100 nM Rosiglitazone (positive control) or serial dilutions (ranging from 10<sup>-3</sup> to 10<sup>-6</sup>) of the different microplastic extracts, showing triglyceride accumulation per well. (c) Graph summarizing NucBlue staining experiments to assess variations in the number of cells, showing DNA content per

well. (d) Graph showing triglyceride accumulation per cell (triglyceride accumulation normalized to DNA content), calculated as the ratio of AdipoRed and NucBlue staining. In b,c,d data are reported as % of control condition (solvent control values were set equal to 100%; horizontal dotted line) and represent the mean  $\pm$  SEM of three independent experiments. \*Indicates dilutions with significant increase over solvent control,  $p < 0.05$ , within each sampling site.

#### 4. Discussion

Despite the growing monitoring of MP presence in the Mediterranean Sea, there is still a substantial lack of information on amount and distribution of MPs in the Adriatic Sea (de Lucia et al., 2014; Gomiero et al., 2018). In this respect, the present study was aimed to provide new insight on the occurrence of MPs in Italian coastal waters of the Central Adriatic Sea and on their capability to adsorb organic pollutants.

Our results revealed that the average amount of MPs (2.65 items/m<sup>3</sup>) estimated in surface offshore waters was similar to that recorded within the Tremiti island Marine Protected Area (Central Adriatic Sea, Italy) in 2017 (2.2 items/m<sup>3</sup>; Mezzelani et al., 2018). Interestingly, this last data was considerably increased with respect to the amount of 0.165 items/m<sup>3</sup> found by (De Lucia et al., 2018) in the same area during 2015. These findings suggest how the Adriatic Sea basin has been characterized by an exponential increase in plastic pollution over the last years. Recently, a range of 0.05 to 4.90 particles/m<sup>3</sup> was reported for quantification of floating plastics (<700  $\mu$ m) in the southern Adriatic Sea by a large scale monitoring study (Suaria et al., 2018). The results of this study also showed that about 51% of the total recorded particles were smaller than 500  $\mu$ m (Gomiero et al., 2018) thus supporting the key role of this size fraction within the plastic pollution affecting the Adriatic Sea (Mezzelani et al., 2018; Zeri et al., 2018). In addition, the high abundance of fragments, as previously observed in other studies (Isobe et al., 2014; Baini et al., 2018), suggests that fragmentation of large plastic objects could represent the main source of MPs in our study area.

340 In the present work, we observed that higher MP abundances were detected in offshore (6 nM)  
341 rather than in inshore (< 1.5 nM) sampling sites. These findings are consistent with the results of  
342 some previous studies which observed prevalence of plastic debris in areas 10-100 Km far from the  
343 coastline, indicating a progressive decrease in the number of plastic items from open sea to coastal  
344 waters (Pedrotti et al., 2016; Bainsi et al., 2018). In contrast, Zeri et al. (2018) have recently found a  
345 statistically significant increases in nearshore ( $\leq 4$  km) microplastic concentration with respect to  
346 that of offshore ( $> 4$  km) Adriatic waters. These contrasting results could be attributed to the  
347 variability in observational conditions which are severely affected by the complexity of the study  
348 area circulation patterns, weather conditions and the heterogeneity in sources of marine litter.

349 A great body of studies is nowadays starting to focus on the potential toxicity of MPs and to  
350 investigate their role as vector of HOCs (Endo et al., 2005; Teuten et al., 2007; Bakir et al., 2014;  
351 Chen et al., 2019; Tang et al., 2020). In this study, the overall evaluation of pollutants carried on  
352 MPs provided a clear separation between in- and offshore waters, highlighting the highest  
353 abundance of these chemicals in the former. This finding is likely due to adsorption mechanisms  
354 from the surrounding environment, because these chemicals, especially PAHs and PCBs, show  
355 affinity with plastic particles (Teuten et al., 2009) and the tendency to accumulate in the organic  
356 phase of sediments (Leggett and Parker, 1994).

357 Among the great variety of organic pollutants which have been isolated on the surface of  
358 microplastics, PHAs (Teuten et al., 2007), PCBs (Endo et al., 2005) and pesticides (Bakir et al.,  
359 2014) represent the most common categories of contaminants. Indeed, PAHs are ubiquitous  
360 pollutants in waters, sediments and marine organisms from the Italian coasts of the Adriatic Sea  
361 (Cocci et al., 2017; Cocci et al., 2018; Frapiccini et al., 2018; Bajt et al., 2019). Due to their strong  
362 ability to bind to plastic particles (Liu et al., 2016), it's not surprising that PAHs are the most  
363 abundant contaminants found on the surface of MPs collected from our study area. The measured  
364 concentration of  $\Sigma$ PAHs is consistent with that found on plastic pellets (166.3 to 2,781 ng/g plastic)  
365 or fragments (85.9 to 2,841 ng/g plastic) beached along Crete shore (Karkanorachaki et al., 2018).

366 Additionally, the levels of PAHs were found to range from 35.1 to 17,023.6 ng/g in marine plastic  
367 debris from Canary Island beaches (Camacho et al., 2019) and from 3400 to 120,000 ng/g in  
368 microplastics from Chinese surface waters (Mai et al., 2018). Our findings are also comparable to  
369 the PAH concentrations (*i.e.* 136.3 to 1586.9  $\mu\text{g/kg}$  plastic and of 397.6 to 2384.2  $\mu\text{g/kg}$  plastic)  
370 recently found in microplastics from two beaches in China (Zhang et al., 2015). Low-ring PAHs  
371 were the most prominent components especially in inshore MPs thus suggesting that petroleum-  
372 derived contaminants are likely to be the main sources of MP contamination. Of the sixteen PAHs,  
373 Acenaphthylene (Acl) showed the highest abundance in each analyzed samples; interestingly, this  
374 finding parallels the highest Acl concentration found in surface coastal waters by our previous  
375 survey (Cocci et al., 2017).

376 PCBs were one of the most abundant group observed in our study showing higher concentrations in  
377 microplastics sampled from coastal areas than in those from offshore waters. These levels are  
378 substantially in line with the measures (*i.e.* 0.9 to 2285.8 ng/g plastic particles) recorded in plastic  
379 pellets and micro fragments beached along the coasts of Canary Islands (Camacho et al., 2019).  
380 PCB concentrations in coastal surface waters of the central Western Adriatic Sea were found to  
381 vary from 6,260 ng/L in areas interested by an intense maritime traffic to 10,650 ng/L in sea basin  
382 affected by riverine runoffs (Cocci et al., 2017). Interestingly, we observed similar patterns in PCB  
383 congener distribution between waters and plastics with highest levels of PCB 52 and 138 (Cocci et  
384 al., 2017). On the other hand, PCB congener 101 and 110 were exclusively identified in  
385 microplastics from the offshore sites or from the harbor, and no detectable concentrations were  
386 found in inshore samples.

387 Total concentrations of OCs (OCs1 and OCs2) were 130.62 ng/g plastic, 19.20 and 59.25 ng/g  
388 plastic in MP from coastal-, offshore waters and harbor, respectively. In the context of pesticides,  
389 there is considerable evidence that OCs are chemically stable under environmental condition, and  
390 their residues can be found in the water even years later then their ban (Breivik et al., 2004). Among  
391 OCs1, we observed that four of them ( $\alpha$ -HCH,  $\beta$ -HCH,  $\gamma$ -HCH and HCB) were found in the

analyzed samples with  $\alpha$ -HCH and  $\beta$ -HCH present in each sampling site. The detected concentration of OC1s (*e.g.* HCB) is consistent with levels found in pellets and plastic fragments from Camacho et al. (2018) but higher (*e.g.*  $\alpha$ -HCH) than values measured in pellets sampled from beaches in China (Zhang et al., 2015). The  $\alpha$ -HCH/ $\gamma$ -HCH ratio was higher than 1, which means that there has been no major recent input of HCHs and that the observed values are due to their persistence in the marine environment. After all, isomeric mixture of HCHs has been used as pesticides for a long time. Recent studies have shown higher concentrations of  $\beta$ -HCH with respect to the other isomers in tissues of mollusks collected at different areas including the Eastern Adriatic (Wang et al., 2008; Kljakovic-Gaspic et al., 2010). These findings are interesting because support the role of environmental conditions in determining the different behaviors of HCH contaminations in different matrices.

Regarding OCs2, 2,4-DDE was found at each sampling site while very high concentration of 4,4-DDD were detected exclusively in the harbor. In addition, high levels of 4,4-DDT were measured in pooled samples from offshore waters. The overall analysis of DDT input in terms of ratio of (DDD + DDE)/DDT (Hitch and Day, 1992) shows a value of  $>0.5$  suggesting presence of DDT residues in the environment. It is not surprising that we found similar values since DDT in the environment is mostly a result of both past production and use (ATSDR, 2002).

OPs are the major representative groups of pesticides found on MPs. In particular, the Chlorpyrifos pesticide was carried by MPs sampled at each site and was found to show the highest concentration among OPs. This is an innovative finding due to the classification of chlorpyrifos as emerging contaminant but, at the same time, it is not surprising because Camacho and coworkers (2019) have recently found this pesticide bound to microplastics collected along the Canary Islands beaches. Chlorpyrifos, malathion and pirimiphos-methyl have been extensively used in agriculture during the last decade and therefore have been detected in biotic and abiotic marine compartments (Henriquez-Hernandez et al., 2017; Moreno-Gonzalez and Leon, 2017). In Italy, chlorpyrifos is largely employed on vineyards and despite the low persistence in the environment occasional



contamination of surface water can be related to point-source pollution (Carter and Capri, 2004; Capri et al., 2005). In addition to these findings, starting in 2007, a particularly interesting massive use of chlorpyrifos was made as part of the integrated pest management (IPM) strategy for *R. ferrugineus* control in the Marche region (Central-Eastern Italy) (Nardi et al., 2011).

Several recent reports have provided evidences that marine microplastics can work as contaminants vectors and that their leachates of chemical pollutants exert possible adverse effects on health (Franzellitti et al., 2019). Although these studies have investigated the activation of both aryl hydrocarbon-receptor (AhR)- and estrogen receptor (ER)-signal transduction pathways, little research has been carried out to investigate the role of MP-associated micropollutants in altering metabolic functions. In this regard, recent evidence proves that environmental pollutants can affect lipid homeostasis promoting obesogenic effects (Cocci et al., 2015; Palermo et al., 2016; Cocci et al., 2017; Cocci et al., 2019). The present study thus aimed to investigate the potential adipogenic properties of MP extracts by using 3T3-L1 adipocytes, a long-established *in vitro* model to assess adipocyte differentiation. Indeed, our results revealed that the pollutants extracted from microplastics recovered in different sites of the Central Adriatic Sea are able to trigger obesogenic effects *in vitro*. Comparatively, it is not possible to define the contribution of each toxicant to the observed effect, because not all components present in the extract may induce the effect chosen for analysis. We can, however, suggest a modulatory influence of pollutants on the effects of other chemicals (found in the extract) that cannot be predicted by adopting the concept of additivity (Kortenkamp, 2007). Our findings are in line with previous observations reporting obesogenic effects of persistent organic pollutants including PCBs, PAHs and OC pesticides (Lee et al., 2012; Rundle et al., 2012; Ferrante et al., 2014). *In vitro* data have demonstrated the obesogenic nature of some PCBs which are likely to modulate adipogenesis via AhR-mediated mechanisms (Arsenescu et al., 2008). Interestingly, two of the PCBs found in our plastic extracts, PCB101 and 138, were able to promote adipogenesis in 3T3-L1 cells (Ferrante et al., 2014; Kim et al., 2017). In particular, these studies suggest that the obesogenicity of PCBs seems to be congener specific.



444 It has also been reported that prenatal exposure to PAH mixture can be associated to obesity  
445 (Rundle et al., 2012). This finding was further confirmed following postnatal exposure to  
446 benzo(a)pyrene that was found to induce similar increases in fat mass (Irigaray et al., 2006). Also  
447 OC pesticides have been suggested as a possible causative agents of obesity due their role in  
448 modulating adipogenesis and lipid metabolism (Lee et al., 2012; Rosenbaum et al., 2017). Previous  
449 *in vitro* data found that both p,p'-DDT and DDE significantly enhanced differentiation of 3T3-L1  
450 preadipocytes (Moreno-Aliaga and Matsumura, 2002; Mangum et al., 2015). For  $\beta$ -HCH, one of the  
451 most common OCs<sup>2</sup> found in our plastic extracts, epidemiological data indicate the presence of a  
452 positive correlation between its serum levels and BMI (Jakszyn et al., 2009; Dirinck et al., 2011).

453 It is clear from the previous findings and the results presented here that the metabolic effects  
454 induced by MP-associated contaminants are due to altered expression of key lipogenic  
455 transcriptional factors such as the PPAR $\gamma$ . Indeed, the signaling cascades that provoke adipogenesis  
456 and lipid accumulation in adipocytes is mediated by the induction of PPAR $\gamma$  (Tontonoz and  
457 Spiegelman, 2008). Most of the chemicals found in the present work have demonstrated the ability  
458 to interact with nuclear receptors acting as agonist of the different PPAR isoforms in both *in vitro*  
459 and *in vivo* models. Previous data from our lab found that exposure to environmentally relevant  
460 doses of several plasticizers induce PPAR $\gamma$ -mediated lipid accumulation in 3T3-L1 adipocytes  
461 (Pomatto et al., 2018). Besides PPAR $\gamma$ , the other PPARs play also a role in tissue fatty acid  
462 metabolism. For example, Cocci et al. (2017) observed an upregulation of PPAR $\alpha$  in fish  
463 hepatocytes exposed to seawater organic extracts from PAHs and PCBs contaminated areas of  
464 Adriatic Sea. Adeogun et al. (2016) found similar effects demonstrating a correlation between the  
465 chemical levels of PAHs and PCBs detected in *Tilapia* ssp sampled from African polluted rivers  
466 and the expression levels of PPARs genes. Therefore, PPAR $\gamma$  seems to be the principal driver  
467 of environmental pollutant-induced adipogenic effects and epigenetic mechanisms are most likely  
468 to be responsible for this action.

Although the present work doesn't investigate the potential transfer of MP-associated contaminants to marine organisms, our results provide further indication of the risk related to litter pollution. Ecotoxicological effects of microplastics have been demonstrated by Avio et al. (2015) showing bioaccumulation of PAHs in mussels exposed to pyrene-coated plastic. Similarly, ingestion of PVC with sorbed triclosan was found to affect feeding behavior or cause mortality in lungworms (Browne et al., 2013). In fish, Rochman et al. (2013; 2014b) showed that long term exposure to plastics coated with a complex mixture of POPs and metals, caused liver toxicity, glycogen depletion, lipidosis, cellular death, tumor promotion, abnormal growth of germ cells in gonads. Overall these findings highlight the importance of combined chemical and physical impacts of microplastics to marine ecosystem.

## Conclusions

In conclusion, this paper reports new preliminary monitoring data, obtained by a standardized sampling protocol, on the quantification of floating microplastics along the coastline in the southern Marche (middle Adriatic). We found a higher abundance of items in the offshore waters (3 - 10 km to the coast) and a general prevalence of fragments as shape category. Land-based sources and riverine inputs are most likely to influence the presence of microplastics, whose distribution depends on the complexity of the study area circulation patterns. There is overall consistency between our findings and those already published for the Adriatic basin. Chemical analysis showed that microplastics, mainly from inshore coastal surface waters, possessed relevant hydrophobic organic contaminants concentrations, including PAHs and pesticides. These findings clearly support the greater sorption ability of marine microplastics. Moreover, *in vitro* 3T3-L1 screening of MP extracts indicated potential metabolic effects resulting in both adipogenesis and lipid uptake/storage. Although our results cannot represent a long-term weather exposure scenario, it is clear from this study and other works that special attention should be paid to the eco-toxicological impact of microplastics, including potential obesogenic effects. It is now evident that combined chemical

and physical effects of microplastics may pose health risks to marine organisms, especially when considering long term exposure in low-flow ecosystems. However, further investigations are needed to elucidate ecotoxicological models in order to ensure a suitable and detailed risk assessment.

#### **Declaration of competing interest**

The authors declare that they have no competing interests.

#### **Acknowledgement**

This work was partially supported by the grant n°EC-172R-18 from the National Geographic Society and by Sky Ocean Rescue.

#### **References**

- Adeogun, A.O., Ibor, O.R., Regoli, F., Arukwe, A., 2016. Peroxisome proliferator-activated receptors and biotransformation responses in relation to condition factor and contaminant burden in tilapia species from Ogun River, Nigeria. *Comparative biochemistry and physiology. Toxicology & pharmacology* : CBP 183-184, 7-19.
- Alimba, C.G., Faggio, C., 2019. Microplastics in the marine environment: Current trends in environmental pollution and mechanisms of toxicological profile. *Environmental toxicology and pharmacology* 68, 61-74.
- Arsenescu, V., Arsenescu, R.I., King, V., Swanson, H., Cassis, L.A., 2008. Polychlorinated biphenyl-77 induces adipocyte differentiation and proinflammatory adipokines and promotes obesity and atherosclerosis. *Environmental health perspectives* 116, 761-768.
- ATSDR, 2002. Agency for Toxic Substances and Disease Registry. Toxicological profile for DDT, DDE, and DDD. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service.

- 520 Avio, C.G., Gorbi, S., Milan, M., Benedetti, M., Fattorini, D., d'Errico, G., Pauletto, M., Bargelloni,  
521 L., Regoli, F., 2015. Pollutants bioavailability and toxicological risk from microplastics to marine  
522 mussels. *Environmental pollution* 198, 211-222.
- 523 Baini, M., Fossi, M.C., Galli, M., Caliani, I., Campani, T., Finoia, M.G., Panti, C., 2018.  
524 Abundance and characterization of microplastics in the coastal waters of Tuscany (Italy): The  
525 application of the MSFD monitoring protocol in the Mediterranean Sea. *Marine pollution bulletin*  
526 133, 543-552.
- 527 Bajt, O., Ramsak, A., Milun, V., Andral, B., Romanelli, G., Scarpato, A., Mitric, M., Kupusovic,  
528 T., Kljajic, Z., Angelidis, M., ulla, A., Galgani, F., 2019. Assessing chemical contamination in the  
529 coastal waters of the Adriatic Sea using active mussel biomonitoring with *Mytilus galloprovincialis*.  
530 *Marine pollution bulletin* 141, 283-298.
- 531 Bakir, A., Rowland, S.J., Thompson, R.C., 2014. Transport of persistent organic pollutants by  
532 microplastics in estuarine conditions. *Estuar Coast Shelf Sci* 140, 14–21.
- 533 Barnes, D.K., Galgani, F., Thompson, R.C., Barlaz, M., 2009. Accumulation and fragmentation of  
534 plastic debris in global environments. *Philosophical transactions of the Royal Society of London.*  
535 *Series B, Biological sciences* 364, 1985-1998.
- 536 Batel, A., Linti, F., Scherer, M., Erdinger, L., Braunbeck, T., 2016. Transfer of benzo[a]pyrene  
537 from microplastics to *Artemia* nauplii and further to zebrafish via a trophic food web experiment:  
538 CYP1A induction and visual tracking of persistent organic pollutants. *Environmental toxicology*  
539 *and chemistry* 35, 1656-1666.
- 540 Besley, A., Vijver, M.G., Behrens, P., Bosker, T., 2017. A standardized method for sampling and  
541 extraction methods for quantifying microplastics in beach sand. *Marine pollution bulletin* 114, 77-  
542 83.
- 543 Breivik, K., Alcock, R., Li, Y.F., Bailey, R.E., Fiedler, H., Pacyna, J.M., 2004. Primary sources of  
544 selected POPs: regional and global scale emission inventories. *Environmental pollution* 128, 3-16.

- 545 Browne, M.A., Crump, P., Niven, S.J., Teuten, E., Tonkin, A., Galloway, T., Thompson, R., 2011.  
 546 Accumulation of microplastic on shorelines worldwide: sources and sinks. *Environmental science &*  
 547 *technology* 45, 9175-9179.
- 548 Browne, M.A., Niven, S.J., Galloway, T.S., Rowland, S.J., Thompson, R.C., 2013. Microplastic  
 549 moves pollutants and additives to worms, reducing functions linked to health and biodiversity.  
 550 *Current biology : CB* 23, 2388-2392.
- 551 Camacho, M., Herrera, A., Gomez, M., Acosta-Dacal, A., Martinez, I., Henriquez-Hernandez, L.A.,  
 552 Luzardo, O.P., 2019. Organic pollutants in marine plastic debris from Canary Islands beaches. *The*  
 553 *Science of the total environment* 662, 22-31.
- 554 Capri, E., Balderacchi, M., Yon, D., Reeves, G., 2005. Deposition and dissipation of chlorpyrifos in  
 555 surface water following vineyard applications in northern Italy. *Environmental toxicology and*  
 556 *chemistry* 24, 852-860.
- 557 Carter, A., Capri, E., 2004. Exposure and effect of chlorpyrifos following use under southern  
 558 European conditions. *Pesticide Outlook* 15, 24-28.
- 559 Chen, Q., Zhang, H., Allgeier, A., Zhou, Q., Ouellet, J.D., Crawford, S.E., Luo, Y., Yang, Y., Shi,  
 560 H., Hollert, H., 2019. Marine microplastics bound dioxin-like chemicals: Model explanation and  
 561 risk assessment. *Journal of hazardous materials* 364, 82-90.
- 562 Cocci, P., Capriotti, M., Mosconi, G., Campanelli, A., Frapiccini, E., Marini, M., Caprioli, G.,  
 563 Sagratini, G., Aretusi, G., Palermo, F.A., 2017. Alterations of gene expression indicating effects on  
 564 estrogen signaling and lipid homeostasis in seabream hepatocytes exposed to extracts of seawater  
 565 sampled from a coastal area of the central Adriatic Sea (Italy). *Marine environmental research* 123,  
 566 25-37.
- 567 Cocci, P., Mosconi, G., Arukwe, A., Mozzicafreddo, M., Angeletti, M., Aretusi, G., Palermo, F.A.,  
 568 2015. Effects of Diisodecyl Phthalate on PPAR:RXR-Dependent Gene Expression Pathways in Sea  
 569 Bream Hepatocytes. *Chemical research in toxicology* 28, 935-947.

- 570 Cocci, P., Mosconi, G., Bracchetti, L., Nalocca, J.M., Frapiccini, E., Marini, M., Caprioli, G.,  
 571 Sagratini, G., Palermo, F.A., 2018. Investigating the potential impact of polycyclic aromatic  
 572 hydrocarbons (PAHs) and polychlorinated biphenyls (PCBs) on gene biomarker expression and  
 573 global DNA methylation in loggerhead sea turtles (*Caretta caretta*) from the Adriatic Sea. The  
 574 Science of the total environment 619-620, 49-57.
- 575 Cocci, P., Mosconi, G., Palermo, F.A., 2019. Changes in expression of microRNA potentially  
 576 targeting key regulators of lipid metabolism in primary gilthead sea bream hepatocytes exposed to  
 577 phthalates or flame retardants. Aquatic toxicology 209, 81-90.
- 578 Cole, M., Galloway, T.S., 2015. Ingestion of Nanoplastics and Microplastics by Pacific Oyster  
 579 Larvae. Environmental science & technology 49, 14625-14632.
- 580 De Lucia, G., Vianello, A., Camedda, A., Vani, D., Tomassetti, P., Coppa, S., Palazzo, L., Amici,  
 581 M., Romanelli, G., Zampetti, G., Cicero, A., Carpentieri, S., Di Vito, S., Matiddi, M., 2018. Sea  
 582 Water Contamination in the Vicinity of the Italian Minor Islands Caused by Microplastic Pollution.  
 583 Water 10, 1108.
- 584 de Lucia, G.A., Caliani, I., Marra, S., Camedda, A., Coppa, S., Alcaro, L., Campani, T., Giannetti,  
 585 M., Coppola, D., Cicero, A.M., Panti, C., Baini, M., Guerranti, C., Marsili, L., Massaro, G., Fossi,  
 586 M.C., Matiddi, M., 2014. Amount and distribution of neustonic micro-plastic off the western  
 587 Sardinian coast (Central-Western Mediterranean Sea). Marine environmental research 100, 10-16.
- 588 Dirinck, E., Jorens, P.G., Covaci, A., Geens, T., Roosens, L., Neels, H., Mertens, I., Van Gaal, L.,  
 589 2011. Obesity and persistent organic pollutants: possible obesogenic effect of organochlorine  
 590 pesticides and polychlorinated biphenyls. Obesity 19, 709-714.
- 591 Endo, S., Takizawa, R., Okuda, K., Takada, H., Chiba, K., Kanehiro, H., Ogi, H., Yamashita, R.,  
 592 Date, T., 2005. Concentration of polychlorinated biphenyls (PCBs) in beached resin pellets:  
 593 variability among individual particles and regional differences. Marine pollution bulletin 50, 1103-  
 594 1114.

- Engler, R.E., 2012. The complex interaction between marine debris and toxic chemicals in the ocean. *Environmental science & technology* 46, 12302-12315.
- Ferrante, M.C., Amero, P., Santoro, A., Monnolo, A., Simeoli, R., Di Guida, F., Mattace Raso, G., Meli, R., 2014. Polychlorinated biphenyls (PCB 101, PCB 153 and PCB 180) alter leptin signaling and lipid metabolism in differentiated 3T3-L1 adipocytes. *Toxicology and applied pharmacology* 279, 401-408.
- Fossi, M.C., Coppola, D., Baini, M., Giannetti, M., Guerranti, C., Marsili, L., Panti, C., de Sabata, E., Clo, S., 2014. Large filter feeding marine organisms as indicators of microplastic in the pelagic environment: the case studies of the Mediterranean basking shark (*Cetorhinus maximus*) and fin whale (*Balaenoptera physalus*). *Marine environmental research* 100, 17-24.
- Fossi, M.C., Panti, C., Guerranti, C., Coppola, D., Giannetti, M., Marsili, L., Minutoli, R., 2012. Are baleen whales exposed to the threat of microplastics? A case study of the Mediterranean fin whale (*Balaenoptera physalus*). *Marine pollution bulletin* 64, 2374-2379.
- Franzellitti, S., Canesi, L., Auguste, M., Wathsala, R., Fabbri, E., 2019. Microplastic exposure and effects in aquatic organisms: A physiological perspective. *Environmental toxicology and pharmacology* 68, 37-51.
- Frapiccini, E., Annibaldi, A., Betti, M., Polidori, P., Truzzi, C., Marini, M., 2018. Polycyclic aromatic hydrocarbon (PAH) accumulation in different common sole (*Solea solea*) tissues from the North Adriatic Sea peculiar impacted area. *Marine pollution bulletin* 137, 61-68.
- Gall, S.C., Thompson, R.C., 2015. The impact of debris on marine life. *Marine pollution bulletin* 92, 170-179.
- Gomiero, A., Strafella, P., Fabi, G., 2018. From Macroplastic to Microplastic Litter: Occurrence, Composition, Source Identification and Interaction with Aquatic Organisms. Experiences from the Adriatic Sea, *Plastics in the Environment*. Alessio Gomiero, IntechOpen, DOI: 10.5772/intechopen.81534.

- 620 Grün, F., Blumberg, B., 2009. Endocrine disrupters as obesogens. Molecular and Cellular  
621 Endocrinology 304 (1-2), 19-29.
- 622 Henriquez-Hernandez, L.A., Carreton, E., Camacho, M., Montoya-Alonso, J.A., Boada, L.D.,  
623 Valeron, P.F., Falcon-Cordon, Y., Falcon-Cordon, S., Almeida-Gonzalez, M., Zumbado, M.,  
624 Luzardo, O.P., 2017. The heartworm (*Dirofilaria immitis*) seems to be able to metabolize  
625 organochlorine pesticides and polychlorinated biphenyls: A case-control study in dogs. The Science  
626 of the total environment 575, 1445-1452.
- 627 Hirai, H., Takada, H., Ogata, Y., Yamashita, R., Mizukawa, K., Saha, M., Kwan, C., Moore, C.,  
628 Gray, H., Laursen, D., Zettler, E.R., Farrington, J.W., Reddy, C.M., Peacock, E.E., Ward, M.W.,  
629 2011. Organic micropollutants in marine plastics debris from the open ocean and remote and urban  
630 beaches. Marine pollution bulletin 62, 1683-1692.
- 631 Hitch, R.K., Day, H.R., 1992. Unusual persistence of DDT in some Western USA soils. Bulletin of  
632 environmental contamination and toxicology 48, 259-264.
- 633 Imhof, H.K., Sigl, R., Brauer, E., Feyl, S., Giesemann, P., Klink, S., Leupolz, K., Loder, M.G.,  
634 Loschel, L.A., Missun, J., Muszynski, S., Ramsperger, A.F., Schrank, I., Speck, S., Steibl, S.,  
635 Trotter, B., Winter, I., Laforsch, C., 2017. Spatial and temporal variation of macro-, meso- and  
636 microplastic abundance on a remote coral island of the Maldives, Indian Ocean. Marine pollution  
637 bulletin 116, 340-347.
- 638 Irigaray, P., Ogier, V., Jacquenet, S., Notet, V., Sibille, P., Mejean, L., Bihain, B.E., Yen, F.T.,  
639 2006. Benzo[a]pyrene impairs beta-adrenergic stimulation of adipose tissue lipolysis and causes  
640 weight gain in mice. A novel molecular mechanism of toxicity for a common food pollutant. The  
641 FEBS journal 273, 1362-1372.
- 642 Isobe, A., Kubo, K., Tamura, Y., Kako, S., Nakashima, E., Fujii, N., 2014. Selective transport of  
643 microplastics and mesoplastics by drifting in coastal waters. Marine pollution bulletin 89, 324-330.
- 644 Jakszyn, P., Goni, F., Etxeandia, A., Vives, A., Millan, E., Lopez, R., Amiano, P., Ardanaz, E.,  
645 Barricarte, A., Chirlaque, M.D., Dorronsoro, M., Larranaga, N., Martinez, C., Navarro, C.,



- Rodriguez, L., Sanchez, M.J., Tormo, M.J., Gonzalez, C.A., Agudo, A., 2009. Serum levels of organochlorine pesticides in healthy adults from five regions of Spain. *Chemosphere* 76, 1518-1524.
- Karkanorachaki, K., Kiparissis, S., Kalogerakis, G.C., Yiantzi, E., Psillakis, E., Kalogerakis, N., 2018. Plastic pellets, meso- and microplastics on the coastline of Northern Crete: Distribution and organic pollution. *Marine pollution bulletin* 133, 578-589.
- Kim, H.Y., Kwon, W.Y., Kim, Y.A., Oh, Y.J., Yoo, S.H., Lee, M.H., Bae, J.Y., Kim, J.M., Yoo, Y.H., 2017. Polychlorinated biphenyls exposure-induced insulin resistance is mediated by lipid droplet enlargement through Fsp27. *Archives of toxicology* 91, 2353-2363.
- Kljakovic-Gaspic, Z., Herceg-Romanic, S., Kozul, D., Veza, J., 2010. Biomonitoring of organochlorine compounds and trace metals along the Eastern Adriatic coast (Croatia) using *Mytilus galloprovincialis*. *Marine pollution bulletin* 60, 1879-1889.
- Kortenkamp, A., 2007. Ten years of mixing cocktails: a review of combination effects of endocrine-disrupting chemicals. *Environmental health perspectives* 115 Suppl 1, 98-105.
- Lee, D.H., Lind, L., Jacobs, D.R., Jr., Salihovic, S., van Bavel, B., Lind, P.M., 2012. Associations of persistent organic pollutants with abdominal obesity in the elderly: The Prospective Investigation of the Vasculature in Uppsala Seniors (PIVUS) study. *Environment international* 40, 170-178.
- Leggett, D.C., Parker, L.V., 1994. Modeling the Equilibrium Partitioning of Organic Contaminants between PTFE, PVC, and Groundwater. *Environmental science & technology* 28, 1229-1233.
- Liu, L., Fokkink, R., Koelmans, A.A., 2016. Sorption of polycyclic aromatic hydrocarbons to polystyrene nanoplastic. *Environmental toxicology and chemistry* 35, 1650-1655.
- Lusher, A.L., Hollman, P.C.H., Mendoza-Hill, J.J., 2017a. Microplastics in fisheries and aquaculture: status of knowledge on their occurrence and implications for aquatic organisms and food safety. *FAO Fisheries and Aquaculture Technical Paper*. No. 615. Rome, Italy.
- Lusher, A.L., Welden, N.A., Sobral, P., Cole, M., 2017b. Sampling, isolating and identifying microplastics ingested by fish and invertebrates. *Analytical Methods* 9(9), 1346-1360.

- 672 Mai, L., Bao, L.J., Shi, L., Liu, L.Y., Zeng, E.Y., 2018. Polycyclic aromatic hydrocarbons affiliated  
673 with microplastics in surface waters of Bohai and Huanghai Seas, China. *Environmental pollution*  
674 241, 834-840.
- 675 Mangum, L.H., Howell, G.E., 3rd, Chambers, J.E., 2015. Exposure to p,p'-DDE enhances  
676 differentiation of 3T3-L1 preadipocytes in a model of sub-optimal differentiation. *Toxicology*  
677 letters 238, 65-71.
- 678 Mato, Y., Isobe, T., Takada, H., Kanehiro, H., Ohtake, C., Kaminuma, T., 2001. Plastic resin pellets  
679 as a transport medium for toxic chemicals in the marine environment. *Environmental science &*  
680 *technology* 35, 318-324.
- 681 Mazurais, D., Ernande, B., Quazuguel, P., Severe, A., Huelvan, C., Madec, L., Mouchel, O.,  
682 Soudant, P., Robbens, J., Huvet, A., Zambonino-Infante, J., 2015. Evaluation of the impact of  
683 polyethylene microbeads ingestion in European sea bass (*Dicentrarchus labrax*) larvae. *Marine*  
684 *environmental research* 112, 78-85.
- 685 Mezzelani, M., Avio, C.G., Nardi, A., Regoli, F., 2018. Microplastic investigation in water and  
686 trophic chain along the Italian coast 06/24/2017-07/15/2017.
- 687 Minambiente, 2018. Programmi di Monitoraggio per la Strategia Marina (Art. 11, D.lgs. 190/2010).  
688 Schede metodologiche per l'attuazione delle Convenzioni stipulate tra Ministero dell'Ambiente e  
689 della Tutela del Territorio e del Mare e Agenzie Regionali per la protezione dell'Ambiente nel  
690 dicembre 2017. Modulo 2: Analisi delle microplastiche, macroplastiche e altri rifiuti flottanti.
- 691 Moreno-Aliaga, M.J., Matsumura, F., 2002. Effects of 1,1,1-trichloro-2,2-bis(p-chlorophenyl)-  
692 ethane (p,p'-DDT) on 3T3-L1 and 3T3-F442A adipocyte differentiation. *Biochemical*  
693 *pharmacology* 63, 997-1007.
- 694 Moreno-Gonzalez, R., Leon, V.M., 2017. Presence and distribution of current-use pesticides in  
695 surface marine sediments from a Mediterranean coastal lagoon (SE Spain). *Environmental science*  
696 *and pollution research international* 24, 8033-8048.

- MSFD, 2013. Technical Subgroup on Marine Litter. Guidance on Monitoring of Marine Litter in European Seas. Joint Research Centre of the European Commission. Scientific and policy reports Report EUR 26113 EN.
- Munari, C., Scoponi, M., Mistri, M., 2017. Plastic debris in the Mediterranean Sea: Types, occurrence and distribution along Adriatic shorelines. *Waste management* 67, 385-391.
- Nardi, S., Ricci, E., Lozzi, R., Marozzi, F., Ladurner, E., Chiabrando, F., Granchelli, L., Verdolini, E., Isidoro, N., Riolo, P., 2011. Control of *Rhynchophorus ferrugineus* (Olivier, 1790) according to EU Decision 2007/ 365/EC in the Marche region (Central-Eastern Italy). *Bulletin OEPP/EPPO Bulletin* 41, 103-115.
- Ogata, Y., Takada, H., Mizukawa, K., Hirai, H., Iwasa, S., Endo, S., Mato, Y., Saha, M., Okuda, K., Nakashima, A., Murakami, M., Zurcher, N., Booyatumanondo, R., Zakaria, M.P., Dung le, Q., Gordon, M., Miguez, C., Suzuki, S., Moore, C., Karapanagioti, H.K., Weerts, S., McClurg, T., Burre, E., Smith, W., Van Velkenburg, M., Lang, J.S., Lang, R.C., Laursen, D., Danner, B., Stewardson, N., Thompson, R.C., 2009. International Pellet Watch: global monitoring of persistent organic pollutants (POPs) in coastal waters. 1. Initial phase data on PCBs, DDTs, and HCHs. *Marine pollution bulletin* 58, 1437-1446.
- Palermo, F.A., Cocci, P., Mozzicafreddo, M., Arukwe, A., Angeletti, M., Aretusi, G., Mosconi, G., 2016. Tri-m-cresyl phosphate and PPAR/LXR interactions in seabream hepatocytes: revealed by computational modeling (docking) and transcriptional regulation of signaling pathways. *Toxicology research* 5, 471-481.
- Pedrotti, M.L., Petit, S., Elineau, A., Bruzard, S., Crebassa, J.C., Dumontet, B., Marti, E., Gorsky, G., Cozar, A., 2016. Changes in the Floating Plastic Pollution of the Mediterranean Sea in Relation to the Distance to Land. *PloS one* 11, e0161581.
- Pellini, G., Gomiero, A., Fortibuoni, T., Ferra, C., Grati, F., Tasseti, A.N., Polidori, P., Fabi, G., Scarcella, G., 2018. Characterization of microplastic litter in the gastrointestinal tract of *Solea solea* from the Adriatic Sea. *Environmental pollution* 234, 943-952.

- Pomatto, V., Cottone, E., Cocci, P., Mozzicafreddo, M., Mosconi, G., Nelson, E.R., Palermo, F.A., Bovolín, P., 2018. Plasticizers used in food-contact materials affect adipogenesis in 3T3-L1 cells. *The Journal of steroid biochemistry and molecular biology* 178, 322-332.
- Rios, L.M., Jones, P.R., Moore, C., Narayan, U.V., 2010. Quantitation of persistent organic pollutants adsorbed on plastic debris from the Northern Pacific Gyre's "eastern garbage patch". *Journal of environmental monitoring : JEM* 12, 2226-2236.
- Rochman, C.M., Hentschel, B.T., Teh, S.J., 2014a. Long-term sorption of metals is similar among plastic types: implications for plastic debris in aquatic environments. *PloS one* 9, e85433.
- Rochman, C.M., Hoh, E., Kurobe, T., Teh, S.J., 2013. Ingested plastic transfers hazardous chemicals to fish and induces hepatic stress. *Scientific reports* 3, 3263.
- Rochman, C.M., Kurobe, T., Flores, I., Teh, S.J., 2014b. Early warning signs of endocrine disruption in adult fish from the ingestion of polyethylene with and without sorbed chemical pollutants from the marine environment. *The Science of the total environment* 493, 656-661.
- Rosenbaum, P.F., Weinstock, R.S., Silverstone, A.E., Sjodin, A., Pavuk, M., 2017. Metabolic syndrome is associated with exposure to organochlorine pesticides in Anniston, AL, United States. *Environment international* 108, 11-21.
- Rundle, A., Hoepner, L., Hassoun, A., Oberfield, S., Freyer, G., Holmes, D., Reyes, M., Quinn, J., Camann, D., Perera, F., Whyatt, R., 2012. Association of childhood obesity with maternal exposure to ambient air polycyclic aromatic hydrocarbons during pregnancy. *American journal of epidemiology* 175, 1163-1172.
- Shim, W.J., Hong, S.H., Eo, S., 2018. Marine Microplastics: Abundance, Distribution, and Composition. *Microplastic Contamination in Aquatic Environments. An Emerging Matter of Environmental Urgency* 1-26.

- 747 Suaria, G., Avio, C.G., Mineo, A., Lattin, G.L., Magaldi, M.G., Belmonte, G., Moore, C.J., Regoli,  
 748 F., Aliani, S., 2016. The Mediterranean Plastic Soup: synthetic polymers in Mediterranean surface  
 749 waters. *Scientific reports* 6, 37551.
- 750 Suaria, G., Avio, C.G., Regoli, F., Aliani, S., 2018. Sub-basin scale heterogeneity in the polymeric  
 751 composition of floating microplastics in the Mediterranean Sea. In *Proceedings of the International*  
 752 *Conference on Microplastic Pollution in the Mediterranean Sea* (pp. 1-7). .
- 753 Tang, Y., Rong, J., Guan, X., Zha, S., Shi, W., Han, Y., Du, X., Wu, F., Huang, W., Liu, G., 2020.  
 754 Immunotoxicity of microplastics and two persistent organic pollutants alone or in combination to a  
 755 bivalve species. *Environmental pollution* 258, 113845.
- 756 Teuten, E.L., Rowland, S.J., Galloway, T.S., Thompson, R.C., 2007. Potential for plastics to  
 757 transport hydrophobic contaminants. *Environmental science & technology* 41, 7759-7764.
- 758 Teuten, E.L., Saquing, J.M., Knappe, D.R., Barlaz, M.A., Jonsson, S., Bjorn, A., Rowland, S.J.,  
 759 Thompson, R.C., Galloway, T.S., Yamashita, R., Ochi, D., Watanuki, Y., Moore, C., Viet, P.H.,  
 760 Tana, T.S., Prudente, M., Boonyatumanond, R., Zakaria, M.P., Akkhavong, K., Ogata, Y., Hirai,  
 761 H., Iwasa, S., Mizukawa, K., Hagino, Y., Imamura, A., Saha, M., Takada, H., 2009. Transport and  
 762 release of chemicals from plastics to the environment and to wildlife. *Philosophical transactions of*  
 763 *the Royal Society of London. Series B, Biological sciences* 364, 2027-2045.
- 764 Tontonoz, P., Spiegelman, B.M., 2008. Fat and beyond: the diverse biology of PPARgamma.  
 765 *Annual review of biochemistry* 77, 289-312.
- 766 UNEP, 2016. Marine plastic debris and microplastics - Global lessons and research to inspire action  
 767 and guide policy change. United Nations Environment Programme, Nairobi.
- 768 Wang, Y., Wang, T., Li, A., Fu, J., Wang, P., Zhang, Q., Jiang, G., 2008. Selection of bioindicators  
 769 of polybrominated diphenyl ethers, polychlorinated biphenyls, and organochlorine pesticides in  
 770 mollusks in the Chinese Bohai Sea. *Environmental science & technology* 42, 7159-7165.
- 771 Zeri, C., Adamopoulou, A., Bojanic Varezic, D., Fortibuoni, T., Kovac Virsek, M., Krzan, A.,  
 772 Mandic, M., Mazziotti, C., Palatinus, A., Peterlin, M., Prvan, M., Ronchi, F., Siljic, J., Tutman, P.,

- 773 Vlachogianni, T., 2018. Floating plastics in Adriatic waters (Mediterranean Sea): From the macro-  
774 to the micro-scale. *Marine pollution bulletin* 136, 341-350.
- 775 Zhang, W., Ma, X., Zhang, Z., Wang, Y., Wang, J., Wang, J., Ma, D., 2015. Persistent organic  
776 pollutants carried on plastic resin pellets from two beaches in China. *Marine pollution bulletin* 99,  
777 28-34.
- 778
- 779

## Highlights

- Floating MPs were collected along the western coasts of the Central Adriatic Sea.
- MPs were found in all sampling stations showing the highest abundance in open waters.
- Chemical analysis showed relevant concentrations of pollutants onto MP surfaces.
- MP extracts induced triglyceride accumulation in 3T3-L1 preadipocytes.
- Our results provide further support for the eco-toxicological impact of MPs.

**Declaration of interests**

☒ The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper. On behalf of all authors, the corresponding author states that there is no conflict of interest.

☐ The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Camerino, May 23<sup>rd</sup>, 2020

Sincerely yours,  
Francesco A. Palermo

